

Florida Gallinule one mile west of this town; it is the only record for the region.—SAMUEL S. DICKEY, *Waynesburg, Pa.*

Some New York City Notes.—I elow I record personal observations of some species either of unusual occurrence, or seen in unusual places in New York City.

REDHEAD (*Marila americana*).—Two were seen on the Jerome Park Reservoir in the Bronx on January 10, 1915.

CANVASBACK (*Marila valisineria*).—On January 10, 1915, I saw seventeen Canvasbacks on the Jerome Park Reservoir. Seven were females. They allowed a close approach, and did not take wing but swam away.

RED-HEADED WOODPECKER (*Melanerpes erythrocephalus*).—I saw one of these birds in partly immature plumage near Riverdale on January 20, 1915.

WHITE-THROATED SPARROW (*Zonotrichia albicollis*).—A flock of eleven of these birds was present in City Hall Park on May 13, 14, 15 and 16, 1914. During the time that I observed them they remained for the most part under some low bushes at the southwest corner of the Park, and seemed quite oblivious to the noise of traffic in Broadway. On two occasions I heard one of them singing.

SCARLET TANAGER (*Piranga erythromelas*).—Two males in full plumage were present in City Hall Park on May 13 and 14, 1914. I saw them on the latter date. Their conspicuous color attracted large crowds and many diverting comments were overheard. The birds were mentioned in the 'Evening Sun' of May 14.

OVENBIRD (*Seiurus aurocapillus*).—I saw three Ovenbirds in City Hall Park on May 13 and 14, 1914. On the latter date they were usually near the Scarlet Tanagers mentioned above. Although they walked about in the center of the grass plots they passed unnoticed by the many persons who were watching the bright colored Tanagers.

VEERY (*Hylocichla fuscescens fuscescens*).—On May 13, 1914, I saw two Veerys in City Hall Park in company with the White-throated Sparrows noted above. They also passed unnoticed.

My experience in bird observation about New York City has been limited to two years, and the occurrence of migratory birds in City Hall Park may not be unusual. However it seemed rather startling to me to find the four last named species in the very heart of the down town section, where thousands of persons are constantly passing and there is an incessant rumble and roar of traffic. It may be of interest to note that the Tanagers were the only ones molested by the hordes of English Sparrows which infest the Park, and that even in their case I observed no serious attacks.—CLIFFORD H. PANGBURN, *Lawrence Park, Bronxville, N. Y.*

Notes from Wisconsin.—**BITTERN** (*Botaurus lentiginosus*).—On July 4, 1914, near Stoughton a Bittern was observed swallowing a snake about twelve inches long. The bird seized it by the head and, after